

Improvement of the foam stability of pilsner reference beer, after addition of hop pectin (from bines and cones), commercial pectin (100%) and montol (100%)




FIG. 1





BEST AVAILABLE COPY

Purity (AUA contents) of the pectin fractions isolated from hops
(bines, cones and waste)




FIG. 2

BEST AVAILABLE COPY

Improvement of the foam stability of pilsner reference beer
after addition of hop pectin from waste and montol (60%)




FIG. 3



BEST AVAILABLE COPY

Improvement of the foam stability of pilsner reference beer
after addition of hop pectin from bines and montol (60%)




FIG. 4



BEST AVAILABLE COPY

Improvement of the foam stability of pilsner reference beer
after addition of hop pectin from cones and montol (60%)





FIG. 5



BEST AVAILABLE COPY

Improvement of the foam stability of pilsner reference beer, after
addition of hop pectin from bines (corrected) and montol (100%)




FIG. 6

BEST AVAILABLE COPY




FIG. 7-1

BEST AVAILABLE COPY





FIG. 7-2

Improvement of the foam stability of pilsner reference beer, after addition of hop pectin from residues of hexane extracts, ethanol extracts and CO₂ extracts. montol (100%), bine pectin, hop cone pectin and commercial pectin (100%)



BEST AVAILABLE COPY




FIG. 8